

POST-OPERATIVE INSTRUCTIONS

 <u>Today Until Bedtime</u>: Alternate 2 Tylenol/acetominophen (2 regular strength or 1 extra strength) and 600mg Advil/ibuprofen (3 regular strength or 1 Motrin) every 2 hours. Substitue Norco (acetominophen + hydrocodone) for Tylenol as needed for pain. Do not exceed 4 grams of Tylenol (acetominophen) over a 24 hour period.

 Next dose:
 _____Tylenol
 _____Tylenol
 _____Ibuprofen, etc., until bedtime

- 2. Next three days: Ibuprofen 600mg, 3 times a day, after meals
- 3. Ice pack outside face of surgical area for the next 2 hours today, alternating 10-15 minutes on, 10-15 minutes off. Warm compress on days 2 & 3 to help reduce swelling.
- 4. **No brushing** of surgical site until instructed by doctor (typically by first or second post-op appointment). Do not pull lip down/out to look at surgical site, this may pull out stitches.
- 5. **Rinsing:** If prescribed, use prescription mouth rinse (Peridex/chlorhexidine) twice daily for 1-2 minutes. Do not eat or drink anything for 30 minutes after rinsing. Avoid vigorous rinsing/swishing for the first 2-3 days following surgery.
- 6. Limit physical activity for the next 48 hours (running, working out, heavy lifting, etc...)
- 7. Periopak (putty): If used, leave in as long as possible then dispose of when it falls off/becomes loose.

8. Healing expectations:

- a. Peak swelling in 2-3 days
- b. Swelling, bruising and black eyes may occur as part of the normal healing process.
- c. Minor bleeding from the surgical areas is normal for the first 24-48 hours after surgery. If bleeding noticably increases, place pressure to site with provided gauze for 10-15 minutes to control. If bleeding persists, place pressure with a moistened tea bag (black tea).
- d. Dissolving stitches will dissolve and unravel leaving threads dangling that can be shortened via cuticle scissors. Do not pull or tug on stitches to remove.
- 9. Diet: Softer foods; eat on opposite side from surgery this week. Avoid hot temperature foods and straws today.
- 10. **Medications:** Take all prescribed medications as instructed. Once started, any prescribed antibiotics should be taken to completion.
- 11. Prescriptions: _
- 12. If you have any questions or problems arise, do not hesitate to call us at 925-398-8903 during office hours. After hours, call Dr. Schellinck at 925-786-6594.